

African Journal of Biotechnology Vol. 11(24), pp. 6535-6541, 22 March, 2012

Available online at <http://www.academicjournals.org/AJB>

DOI: 10.5897/AJB11.2789

ISSN 1684-5315 © 2012 Academic Journals

Full Length Research Paper

Effects of over-expression of allene oxide cyclase on camptothecin production by cell cultures of *Camptotheca acuminata*

Yan Pi¹, Keji Jiang^{1,2}, Juan Lin¹, Ying Cao¹, Qian Wang¹, Ji Yang¹,
Xiaofen Sun¹ and Kexuan Tang^{1,3*}

¹State Key Laboratory of Genetic Engineering, School of Life Sciences, Morgan-Tan International Center for Life Sciences, Fudan-SJTU-Nottingham Plant Biotechnology R&D Center, Fudan University, Shanghai 200433, P. R. China.

²East China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Shanghai 200090, China.

³Plant Biotechnology Research Center, School of Agriculture and Biology, Fudan-SJTU-Nottingham Plant Biotechnology R&D Center, Institute of Systems Biology, Shanghai Jiao Tong University, Shanghai 200030, P. R. China.

Accepted 2 March, 2012

Camptothecin (CPT) is an anticancer and antiviral monoterpene-derived indole alkaloid which can be induced by plant hormone, jasmonates. To improve the production of the pharmaceuticals, the jasmonate biosynthesis related gene allene oxide cyclase from *Camptotheca acuminata* was transferred back into *C. acuminata* using the method of *Agrobacterium*-mediated genetic transformation and over expressed. The results of high performance liquid chromatography (HPLC) analysis indicated that the camptothecin content of transgenic callus was higher than that of non-transgenic callus. The highest camptothecin content in transgenic callus was 3.9310 mg/g DW. However, camptothecin content in both transgenic and non-transgenic calli significantly decreased after further extrinsic methyl-jasmonate's (MeJA's) induction, whereas the content of CPT in transgenic callus was still higher than in non-transgenic one. All the results indicate that endogenous jasmonate's accumulation may be promoted after allene oxide cyclase gene was transformed into *C. acuminata* and over expressed. In this way, jasmonates can affect second metabolism pathway genes' expression and then the camptothecin content improved. Inexpectantly, the mechanism of extrinsic jasmonate to secondary metabolism of *C. acuminata* was different from that of endogenous jasmonates.

Key words: *Agrobacterium tumefaciens*, Allene oxide cyclase, *Camptotheca acuminata*, camptothecin, methyl-jasmonates, transformation.

INTRODUCTION

Camptothecin (CPT), originally isolated from the extract of a traditional Chinese medicinal tree *Camptotheca acuminata* (Nyssaceae) (Wall et al., 1966), is a water-insoluble therapeutically valuable monoterpene-derived indole alkaloid. This compound exhibits an ability to specifically inhibit the cleavage and rejoining reaction of DNA topoisomerase I (Topo I) (Hsiang et al., 1985). At

present, semi-synthetic water-soluble camptothecin analogues, topotecan and irinotecan, are used as clinical anti-tumor agents throughout the world. They are synthesized from natural camptothecin, which is mainly obtained by extraction and purification from *C. acuminata* plant. Due to the present and potential clinical uses of CPT and its derivatives, it is important to develop sustainable and alternative production sources of these compounds due to low content of these compounds in natural plants.

Terpenoid indole alkaloids (TIAs) biosynthesis in *Catharanthus roseus* is one of the best studied elicitor-induced secondary metabolic pathways (Noé et al., 1984;

*Corresponding author. E-mail: kxtang@fudan.edu.cn or kxtang1@yahoo.com. Tel: +86 21 6564 2425. Fax: +86 21 6564 2425.




Figure 1. Schematic maps of transformation plasmid (pCambia1304+CaAOC). 35S-Pro, cauliflower mosaic virus 35S promoter. *hygromycin*, *hygromycin B* phosphotransferase gene. Nos-Ter, 3' termination region from *Agrobacterium nopaline* synthase gene. CaAOC, CaAOC gene. Restriction enzyme (*SpeI* and *BstEII*), indicated by arrow.

van der fits and Memelink, 2000, 2001; Pauw et al., 2004). Jasmonate (JA) and its derivative, methyl-jasmonate (MeJA) were reported to be general inducers of plant secondary metabolite biosynthesis (Memelink et al., 2001). The strictosidine synthase (STR; EC 4.3.3.2) and the tryptophan decarboxylase (TDC; EC 4.1.1.28) are essential and are key enzymes in TIA biosynthesis, which are coordinately regulated by a JA-responsive transcription factor (ORCA3) in *C. roseus*. It has showed that ectopic expression of the transcription factor (ORCA3) could enhance the production of TIAs in cultured cells of *C. roseus* (van der Fits and Memelink, 2000, 2001). Maybe it is also an alternative approach that the key genes in jasmonate biosynthetic pathway are modified by genetic manipulation for the production of valuable secondary metabolites which is induced by jasmonates.

An essential enzyme involved in JA biosynthesis is allene oxide cyclase (AOC; EC 5.3.99.6). The AOC-catalyzed step is regarded as the crucial point in the biosynthesis of JA due to the establishment of the naturally occurring enantiomeric structure of JA (Cenzano et al., 2007; Maucher et al., 2004). It was reported that the overexpression of an heterologous AOC gene from *Hyoscyamus niger* L. in *Nicotiana tabacum* cv. Petit Havana can result in overexpression of nicotine biosynthetic pathway genes and higher yields of nicotine (Jiang et al., 2009). The engineering TIA biosynthesis, as an essential step to ensure the profitable of cell suspensions *in vitro* production systems, will face a major challenge for the next years.

In this study, we attempted to investigate the role of AOC gene from JA biosynthesis in promotion of the CPT biosynthetic pathway. An AOC gene (defined as CaAOC, Database Accession No. AY863428), which has been isolated from *C. acuminata* in previous work was

transformed into *C. acuminata*. The result of RT-PCR showed the CaAOC expression level in transgenic callus was obviously higher than that of non-transgenic callus. The results of high performance liquid chromatography (HPLC) analysis indicated that the camptothecin content of transgenic callus was higher than that of non-transgenic callus. In addition, the effect of extrinsic MeJA on the camptothecin content was evaluated as well.

EXPERIMENTALS

Construction of CaAOC expression vector

The vector pCambia1304 (Cambia, Canberra, Australia) used in this study contains a 1.9 kb GUS expression cassette and a hygromycin B phosphotransferase gene (*hpt*) driven by the CaMV35S promoter. To construct pCambia1304+CaAOC, the GUS expression cassette of pCambia1304 was replaced with the *SpeI*-*BstEII* CaAOC coding sequence amplified with FCaAOCp1304 (5'-GG ACTAGT ATG GCT GCT TCA TCA ACT TC-3', the *SpeI* digestion site is underlined) and RCaAOCp1304 (5'-GA GGTTACC TCA GTC AGT GAA GTT AGG AA-3', the *BstEII* digestion site is underlined), using cDNA of *C. acuminata* (Figure 1). The plasmid was introduced into the disarmed *Agrobacterium tumefaciens* strain EHA105. A positive clone, after confirmation by PCR and enzymatic digestion analysis for the presence of the CaAOC gene, was used for plant transformation.

Plant material and *C. acuminata* transformation

A. tumefaciens EHA105 harboring the vector was grown at 28°C in Luria Bertani (LB) medium (Sambrook et al., 2002) supplemented with 100 µg/ml kanamycin, 40 µg/ml streptomycin, 25 µg/ml rifampicin and 100 µM acetosyringone. For infection, the concentration of *Agrobacterium* suspension in inoculation MS (Murashige and Skoog, 1962) liquid medium with 100 µM acetosyringone was adjusted to an OD₆₀₀ of 0.6. Mature seeds of *C. acuminata* were collected from the plants growing in the campus of




Figure 2. Selection of transgenic cell lines. a, Transformed *Camptotheca acuminata* cell lines in selection medium MS₂. b, Transformants selected in MS₂.

Fudan University, Shanghai, China. The decoated seeds were surface sterilized with 0.1% (w/v) of mercuric chloride for 5.0 min and rinsed thoroughly five times in sterile distilled water. Then, the wounded seed embryos were inoculated with *Agrobacterium* for 5.0 min, blotted twice on sterile filter paper and placed on MS₁ callus induction medium (MS solid medium supplemented with 5 µg/ml naphthalene-acetic acid, 0.5 µg/ml 6-benzyladenine, 0.3 µg/ml 2,4-dichlorophenoxy-acetic acid, and 30 mg/ml of sucrose) for two days as a co-culture stage.

Selection and culture of *C. acuminata* cell lines

After the co-culture stage, the wounded seed embryo pieces was transferred onto selection medium MS₂, the same medium as MS₁ but supplemented with 250 µg/ml carbenicillin (Carb) and 20 µg/ml hygromycin B. After initiation of the independent callus initiated from seed embryo in about three to five weeks (Figure 2), it was maintained as usual in darkness (28°C) and was sub-cultured every four weeks to the same fresh selection medium MS₂. Nine independent transgenic cell lines, maintained in selection medium MS₃ (MS₁ supplemented with 20 µg/ml hygromycin B), were selected for examination.

Molecular analysis of transgenic cell lines

Total genomic DNA was extracted from fresh cell lines by cetyltrimethylammonium bromide (CTAB) method as described (Doyle and Doyle, 1990). PCR primers used for detecting the *hpt* gene were Fhpt (5'-CGA TTT GTG TAC GCC CGA CAG TC-3') and Rhpt (5'-CGA TGT AGG AGG GCG TGG ATA TG-3'). Primers for the detection of transformed *CaAOC* were F35Sp1304 (5'-GCA CAA TCC CAC TAT CCT TCG CAA-3') and RCaAOC (5'-CGT ACA CGT CAC CTT TCT TCT CCG G-3'). Plasmid DNA from the *Agrobacterium* carrying pCAMBIA1304+*CaAOC* was used as a

positive control. For detection of the *hpt* gene and the *CaAOC* gene, the template was denatured at 94°C for 3 min followed by 34 amplification cycles (50 s at 94°C, 50 s at 60°C and 50 s at 72°C) and finally at 72°C for 8 min. Products (10 µl) were analyzed on 1% agarose gel.

MeJA treatment

To study the changes of the transcription of *CaAOC* in transgenic cell lines and non-transgenic cell lines under MeJA treatments, the 15-day-old *C. acuminata* cells, sub-cultured on fresh medium in a new Petri dish, were soaked in a solution of 100 µM MeJA, respectively. The cells were grown at 25 ± 2°C in the dark for 60 min, then each sample was divided into two parts, one was stored -80°C using for total RNA isolation, the other was oven-dried at 60°C in the dark for three days to a constant weight and CPT content was determined.

RT-PCR analysis of *CaAOC* in transgenic cell lines

After MeJA induction, the nine transgenic cell lines, accompanied with control treated by distilled water, were subjected to analysis for the transcription of *CaAOC* using One Step RT-PCR (Takara, Shiga, Japan) with the primers *CaAOC*-RT-F (5'-AAACCTCAGACTCA-CCACCACA -3') and *CaAOC*-RT-R (5'-TTCCACGGTCT-CGTTCTGTT -3'). 18S rRNA was used as internal control with the primers 18S-F (5'-ATG ATA ACT CGA CGG ATC GC-3') and 18S-R (5'-CTT GGA TGT GGT AGC CGT TT-3'). RT-PCR was carried out following the manufacturer's protocol, and the thermal cycle condition used was 30 min at 50°C, 2 min at 94°C, followed by 22 cycles of amplification (30 s at 94°C, 30 s at 60°C and 60 s at 72°C). Products (5 µl) were analyzed on 1% agarose gel.




Figure 3. PCR analysis of the transgenic cell lines. M, DNA size marker DL2000 (100 to 2000 bp). P, plasmid (positive control). W, non-transgenic cell lines (negative control). *CaAOC*, *CaAOC* gene (536 bp). *Hpt*, hygromycin B phosphotransferase gene (812 bp). Arabic numbers indicate the independent transgenic cell lines.

Camptothecin extraction and HPLC analysis

The CPT was extracted from dried samples by supersonic oscillator method as described (Yan et al., 2002). After filtering, the solutions were directly analyzed by HPLC (Model 2690; Waters, Milford, MA, USA) as described (Wiedenfeld et al., 1997).

RESULTS

C. acuminata transformation and molecular analysis

The embryos with papery cotyledons from *C. acuminata* were transformed with *A. tumefaciens* strain EHA105 harboring pCambia1304+*CaAOC*. Transformed cell lines were excised and put onto selection medium MS₂ and maintained in selection medium MS₃. The putative transformants were analyzed by PCR to test the integration of the exogenous gene. In order to avoid the effect of inherent AOC gene in *C. acuminata*, a section of the 35S promoter sequences was used to explore *CaAOC* gene as a forward primer. PCR was performed with the DNA from transformed plants using *hpt* and *CaAOC* gene primers. The products were analyzed on 1% agarose gel, the expected sizes of the amplified fragments were 812 and 536 bp (Figure 3). The untransformed plants did not show any signals.

RT-PCR analysis of *CaAOC* in transgenic *C. acuminata* cells

It was reported that the transcription of AOC gene nearly reached their higher levels at 60 min after treated with MeJA (Agrawal et al., 2003a, b; Stenzel et al., 2003). To examine the changes of *CaAOC* gene expression under extrinsic MeJA treatments, the 15-day-old *C. acuminata* calli were used for RNA isolation after treated for 60 min. The result shows that the transcript of *CaAOC* was induced by 100 μ M MeJA in the nine transgenic calli and non-transgenic control callus (Figure 4). Under extrinsic MeJA induction, the expression of *CaAOC* mRNA in transgenic calli was higher than that in the non-transgenic control.

Camptothecin content analysis in transgenic *C. acuminata* cells

The profiles of camptothecin concentration in transgenic cell lines were determined by HPLC. The capacities of transgenic cell lines to biosynthesize camptothecin are shown in Figure 5. The content of camptothecin in transgenic calli (it ranged from 3.9310 mg/g to 1.0028 mg/g of dry weight) was much higher than that in non-




Figure 4. RT-PCR analysis of the transgenic cell lines after 100 μ M MeJA treatments. *CaAOC*, *CaAOC* gene. *18S*, *18S rRNA*. W, non-transgenic cell lines. Arabic numbers indicate the independent transgenic cell lines. MeJA(-), Treatments without MeJA; MeJA(+), treatments with MeJA.

transgenic control which was nearly detected. The 3.9310 mg/g was higher than those values described in the literature before and was comparable to the results of young flower buds and seeds from *C. acuminata* plant.

DISCUSSION

Increasing evidences indicate that jasmonates can regulate the synthesis of secondary metabolites in many species by an ubiquitous plant defense response (Keinänen et al., 2001; Goossens et al., 2003; Sasaki-Sekimoto et al., 2005; Wang et al., 2010), and the transgenic manipulation of endogenous JA levels in tomato and tobacco proved to be feasible (Jiang et al., 2009; Keinänen et al., 2001; Chen et al., 2006). Our results indicate that *CaAOC* gene had been integrated in the

genome and the transcript level was increased. The CPT accumulation in nine transgenic cell lines was higher than in non-transgenic cell lines. Hairy roots derived from the wild-type strains of *Agrobacterium rhizogenes* have also been developed in *Ophiorrhiza pumila* and *C. acuminata*, the yields of the camptothecin in the two species were of approximately 1 mg/g DW. The CPT content in *C. acuminata* callus in optimization condition was 2.36 mg/g DW. The highest CPT content of the transgenic callus reached 3.9310 mg/g DW in this present research, which was higher than those reported. And the content in organs of the plant seems to be somewhat smaller than that in transgenic calluses but is also in the same range. It could be deduced that the CPT accumulation was associated with increased levels of transcripts that encode *CaAOC* enzyme in the jasmonate pathway. The cloning and overexpression of the JA-responsive




Figure 5. The CPT concentration of transgenic *Camptotheca acuminata* cell lines and non-transgenic cell lines. W, Non-transgenic cell lines. C1 to C9, Nine independent transgenic cell lines. MeJA(-), Treatments without MeJA; MeJA(+), treatments with MeJA.

transcription factors (ORCA) had been shown to regulate the jasmonate-responsive activation of several mono-terpene indole alkaloids-synthesizing genes in cell cultures (Gantet and Memelink, 2002; Vom Endt et al., 2002; Pasquali et al., 2006). The transformation of *CaAOC* in *C. acuminata* might also promote the accumulation of JAs. The transcription factors (ORCA) responded to JAs, and regulated the activation of several CPT biosynthesis pathway genes. Then the CPT production in transgenic callus was enhanced. This might be a way that metabolic engineering and physiological manipulation was used as complementary to improve alkaloid yields.

Extrinsic JAs can elicit accumulation of secondary metabolites in many plants. In this present study, the results show that the transcript of *CaAOC* was induced in the nine transgenic calli and non-transgenic control callus of *C. acuminata* by 100 μ M MeJA (Figure 4), but the CPT accumulation was not simultaneously associated with the increased expression of *CaAOC* in transgenic calli. It was reported that the extrinsic MeJA could significantly enhance the production of secondary metabolites such as the anti-cancer drug paclitaxel (Taxol) in cell suspension cultures of *Taxus* and the TIAs in cells of *C. roseus* (Wang et al., 2010; Ketchum et al., 2003; Magdi and Rob, 2005; Sabah et al., 2007). And MeJA and JA could increase CPT production six and 11 times in *C. acuminata* cells, respectively (Song and Byun, 1998). Our results is not consistent with those cases, but the same as the result previously reported that extrinsic MeJA did not elicit CPT yields in *C. acuminata* cells. What's more, the transgenic lines exhibited significantly diverse CPT concentrations without MeJA treatment, while with MeJA treatment, these lines seemed to perform lower but much closer values in CPT concentration (Figure 5). Herein, possible explanations might lay on that: (1) endogenous MeJA had different affection mechanism on the yields of

TIAs with extrinsic one; (2) in the different time course, MeJA showed different affection mechanism, inhibition or inducement; (3) MeJA had another affection mechanism on the yields of CPT in *C. acuminata* compared with that on the yield of TIAs in *C. roseus* and (4) over accumulation of exo-genous MeJA might have preferentially triggered other secondary metabolism which competes with CPT biosynthesis. On this occasion, all the transgenic lines in MeJA treatment preferred to produce CPT at a restricted and balanced concentration.

Regulation of JA biosynthesis is quite a complex process. It is less understood that the functional role of the subsequent transcriptional up-regulation of JA-biosynthetic enzymes. Here, we offered an alternative to elicitors as a means to enhance the CPT production. The signaling property of JAs is interesting not only for what it reveals about how primary and secondary metabolism is reconfigured in response to stress, but also for practical applications in the production of economically useful compounds.

ACKNOWLEDGEMENTS

The authors thank Mr. Ping Zhang from Chemistry Analysis and Testing Centre in Fudan University for the help with HPLC assay. This work was funded by National Basic Research Program of China (No. 2007CB108805), Shanghai Science and Technology Committee (No. 08391911800), and Research Fund for the Doctoral Program of Higher Education of China (No. 20090071120040).

Abbreviations

AOC, Allene oxide cyclase; **CPT**, camptothecin; **HPLC**,

high performance liquid chromatography; **hpt**, hygromycin B phosphotransferase gene; **JA**, jasmonate; **MeJA**, methyl-jasmonate; **TIAs**, terpenoid indole alkaloids.

REFERENCES

- Agrawal GK, Jwa N-S, Agrawal SK, Tamogami S, Iwahashi H, Rakwal R (2003a). Cloning of novel rice allene oxide cyclase (OsAOC): mRNA expression and comparative analysis with allene oxide synthase (OsAOS) gene provides insight into the transcriptional regulation of octadecanoid pathway biosynthetic genes in rice. *Plant Sci.* 164: 979-992.
- Agrawal GK, Jwa NS, Shibato J, Han O, Iwahashi H, Rakwal R (2003b). Diverse environmental cues transiently regulate OsOPR1 of the "octadecanoid pathway" revealing its importance in rice defense/stress and development. *Biochem. Biophys. Res. Commun.* 310: 1073-1082.
- Cenzano A, Abdala G, Hause B (2007). Cytochemical immuno localization of allene oxide cyclase, a jasmonic acid biosynthetic enzyme, in developing potato stolons. *J. Plant Physiol.* 164: 1449-1456.
- Chen H, Jones AD, Howe GA (2006). Constitutive activation of the jasmonate signaling pathway enhances the production of secondary metabolites in tomato. *FEBS. Lett.* 580: 2540-2546.
- Doyle JJ, Doyle JL (1990). Isolation of plant DNA from fresh tissue. *Focus*, 12: 13-15.
- Gantet P, Memelink J (2002). Transcription factors: tools to engineer the production of pharmacologically active plant metabolites. *Trends Pharmacol. Sci.* 23: 563-569.
- Goossens A, Hakkinen ST, Laakso I, Seppanen-Laakso T, Biondi S, De Sutter V, Lammertyn F, Nuutila AM, Soderlund H, Zabeau M, Inze D, Oksman-Caldentey KM (2003). A functional genomics approach toward the understanding of secondary metabolism in plant cells. *Proc. Natl. Acad. Sci. USA*, 100: 8595-8600.
- Hsiang YH, Hertzberg R, Hecht S, Liu LF (1985). Camptothecin induces protein-linked DNA breaks via mammalian DNA topoisomerase I. *J. Biol. Chem.* 260: 14873-14878.
- Jiang KJ, Pi Y, Hou R, Jiang LL, Sun XF, Tang K (2009). Promotion of nicotine biosynthesis in transgenic tobaccos by overexpression allene oxide cyclase from *Hyoscyamus niger*. *Planta*, 229: 1057-1063.
- Keinänen M, Oldham NJ, Baldwin IT (2001). Rapid HPLC screening of jasmonate-induced increases in tobacco alkaloids, phenolics, and diterpene glycosides in *Nicotiana attenuate*. *J. Agric. Food. Chem.* 49: 3553-3558.
- Ketchum RE, Rithner CD, Qiu D, Kim YS, Williams RM, Croteau RB (2003). Taxus metabolomics: methyl jasmonate preferentially induces production of taxoids oxygenated at C-13 in *Taxus media* cell cultures. *Phytochemistry*, 62: 901-909.
- Magdi E-S, Rob V (2005). Methyljasmonate accelerates catabolism of monoterpenoid indole alkaloids in *Catharanthus roseus* during leaf processing. *Fitoterapia*. 76: 83-90.
- Maucher H, Stenzel I, Miersch O, Stein N, Prasad M, Zierold U, Schweizer P, Dorer C, Hause B, Wasternack C (2004). The allene oxide cyclase of barley (*Hordeum vulgare* L.) – cloning and organ-specific expression. *Phytochemistry*. 2004, 65: 801-811.
- Memelink J, Verpoorte R, Kijne JW (2001). ORCAization of jasmonate-responsive gene expression in alkaloid metabolism. *Trends. Plant Sci.* 6: 212-219.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plantarum*. 15: 473-497.
- Noé W, Mollenschott C, Berlin J (1984). Tryptophan decarboxylase from *Catharanthus roseus* cell suspension cultures: purification, molecular and kinetic data of the homogenous protein. *Plant Mol. Biol.* 3: 281-288.
- Pasquali G, Porto DD, Fett-Neto AG (2006). Metabolic engineering of cell cultures versus whole plant complexity in production of bioactive monoterpene indole alkaloids: recent progress related to old dilemma. *J. Biosci. Bioeng.* 101: 287-296.
- Pauw B, Hilliou FA, Martin VS, Chatel G, de Wolf CJ, Champion A, Pré M, van Duijn B, Kijne JW, van der Fits L, Memelink J (2004). Zinc finger proteins act as transcriptional repressors of alkaloid biosynthesis genes in *Catharanthus roseus*. *J. Biol. Chem.* 279: 52940-52948.
- Sabah H, Vincent C, Nathalie G-G, Pascal G (2007). Regulation of the terpene moiety biosynthesis of *Catharanthus roseus* terpene indole alkaloids. *Phytochem. Rev.* 6: 341-351.
- Sambrook J, Fritsch EF, Maniatis T (2002). Molecular cloning: a laboratory manual, 3rd edn. Cold Spring Harbor Laboratory, Cold Spring Harbor. pp. 1595-1595.
- Sasaki-Sekimoto Y, Taki N, Obayashi T, Aono M, Matsumoto F, Sakurai N, Suzuki H, Hirai MY, Noji M, Saito K, Masuda T, Takamiya K, Shibata D, Ohta H (2005). Coordinated activation of metabolic pathways for antioxidants and defence compounds by jasmonates and their roles in stress tolerance in *Arabidopsis*. *Plant J.* 44: 653-668.
- Song SH, Byun SY (1998). Characterization of cell growth and camptothecin production in cell cultures of *Camptotheca acuminata*. *J. Microbiol. Biotechnol.* 8: 631-638.
- Stenzel I, Hause B, Miersch O, Kurz T, Maucher H, Weichert H, Ziegler J, Feussner I, Wasternack C (2003). Jasmonate biosynthesis and the allene oxide cyclase family of *Arabidopsis thaliana*. *Plant Mol. Biol.* 51: 895-911.
- van der Fits L, Memelink J (2000). ORCA3, a jasmonateresponsive transcriptional regulator of plant primary and secondary metabolism. *Science*, 289: 295-297.
- van der Fits L, Memelink J (2001). The jasmonate-inducible AP2/ERF-domain transcription factor ORCA3 activates gene expression via interaction with a jasmonate-responsive promoter element. *Plant J.* 25: 43-53.
- Vom Endt D, Kijne JW, Memelink J (2002). Transcription factors controlling plant secondary metabolism: what regulates the regulators? *Phytochemistry*, 61: 107-114.
- Wall ME, Wani MC, Cook CE, Palmer AH, McPhail AT, Sim GA (1966). Plant antitumor agents. I. The isolation and structure of camptothecin, a novel alkaloidal leukemia and tumor inhibition from *Camptotheca acuminata*. *J. Am. Chem. Soc.* 88: 3888-3890.
- Wang Q, Yuan F, Pan QF, Li MY, Wang GF, Zhao JY, Tang KX (2010). Isolation and functional analysis of the *Catharanthus roseus* deacetylindoline-4-O-acetyltransferase gene promoter. *Plant Cell. Rep.* 29: 185-192.
- Wiedenfeld H, Furmanowa M, Roeder E, Guzewska J, Gustowski W (1997). Camptothecin and 10-hydroxycamptothecin in callus and plantlets of *Camptotheca acuminata*. *Plant Cell. Tissue Org. Cult.* 49: 213-218.
- Yan XF, Wang Y, Yu T, Zhang YH, Yin LJ (2002). Determination of camptothecin in leaves of *Camptotheca Acuminata* Decne by HPLC. *J. Instrumental Anal.* 21: 15-17 (in Chinese with English abstract).